

Mandatory Reflex Testing

Approved by MEC November 2021

INITIAL TEST and RESULT	CONFIRMATION TESTING/ ADDITIONAL WORKUP
All newly diagnosed patients with Acute Myeloid Leukemia, unless otherwise specified by physician. (Bone Marrow or Whole Blood)	Heme Molecular Sequence Analysis
aPTT with no endpoint detected or error code on the coagulation analyzer that cannot be resolved.	Unfractionated Heparin
aPTT Mixing Study	APTT and, if the APTT Mixing Study does not correct, order a Lupus Anticoagulant Screen (LA1)
aPTT Mixing Studies (Test Code 4120) greater than or equal to 6 seconds above the normal range.	Heparin Neutralization (Policy 6679) to rule out anticoagulant: <ul style="list-style-type: none"> • If aPTT corrects to normal, then a Mixing Study is not indicated. • If aPTT remains elevated, then Unfractionated Heparin Level (Test Code 8101) assay is ordered. <ul style="list-style-type: none"> • If Unfractionated Heparin Level is greater than 1.0 U/mL, then a Mixing Study is not indicated.
Anaerobic Culture	Aerobic culture on all orders that do not already have an aerobic culture ordered on the same specimen.
Antibody Screens	ABO/RH
AntiNuclear Antibody (ANA) EIA Screen	AntiNuclear Antibody (ANA) Hep-2 Substrate with Reflex positive
AntiNuclear Antibody (ANA) Hep-2 Substrate with Reflex positive	AntiNuclear Antibody (ANA) Titer and Pattern
Blood Culture (Test Codes 8894 & 8896); if positive for growth of bacteria	<ul style="list-style-type: none"> • Bacterial identification will be performed if growth occurs any bottle. An antibiotic susceptibility will be performed on all pathogenic isolates.
All newly diagnosed patients with B-Cell Lymphomas with features concerning possible lymphoplasmacytic, unless otherwise specified by physician	<ul style="list-style-type: none"> • MYD88 L265P mutation testing on bone marrow aspirate
All patients that are identified with hemoglobinopathy including: <ul style="list-style-type: none"> • hgSS • hgSC • beta thalassemia And have not been transfused in the last 3 months.	Serological C, E, and K antigen typing
Women of childbearing age identified as RHD variants or "weak D phenotypes" via serological testing with no previous RHD genotyping on file.	Molecular RHD Genotyping
All patients with Invasive ductal, lobular, mixed ductal/lobular, mammary NOS, or micrometastatic lymph node breast carcinoma meeting the following criteria: <ul style="list-style-type: none"> • Age less than 70 years • Not pure tubular, mucinous, or colloid carcinoma (grade 1 special subtypes with good prognosis) • Tumor is pathologic stage pT1b, T1c, T2, or T3 • Tumor is not stage T4 • Tumor is pathologic stage N0 or N1mi • Tumor is ER positive and Her2/neu negative • Not post-treatment (y), recurrent tumor (r) or with known distant metastatic tumor (M1) 	Send for Oncotype DX testing: <ol style="list-style-type: none"> 1) If multifocal, and meet above criteria 2) . Perform Oncotype on largest primary tumor (if same histology) 3) Perform Oncotype on up to three primary tumors (if different histology), indicate order of testing to Oncotype (as they will stop further testing if high recurrence score is result) 4) If bilateral, perform Oncotype on both sides.

INITIAL TEST and RESULT	CONFIRMATION TESTING/ ADDITIONAL WORKUP
	5) If high risk tumor: stage (T4, N1b+), ER-, or Her2 positive concurrent cancer that would not reflex, do not reflect test lower risk tumor(s)
All newly diagnosed or previously untested Adult Granulosa Cell Tumors, Serous Boderline Tumors, and Low Grade Serous Carcinomas	IHC for Estrogen and Progesterone receptors
Bone Marrow samples with clinical indications for possible need to perform FISH Multiple Myeloma Panel	Sort CD138 to purify (concentrate) sample and hold for possible FISH Multiple Myeloma Panel
Diagnostic sample of B-lymphoblastic leukemia (B-ALL) and B-cell non-Hodgkin lymphoma with anti CD-19 therapy	Flow cytometry Blinatumomab tube (anti-CD19 therapy tubes)
CBC w/man diff - will reflex to CBC w/out diff "if" the WBC is less than 0.4.	CBC w/out differential
CBC w/man diff - will reflex to CBC w/out diff "if" requested within in 24 hours.	CBC w/out differential
Flow cytometry testing	<ol style="list-style-type: none"> 1. CBC w diff on all orders that do not already have a CBC w diff ordered on patient on the same date and specimen is less than 10 hours old. 2. And Flow cytometry is unable to get WBC and automated differential.
<p>Celiac Disease Cascade</p> <p>Celiac TTG Ab IgA and Celiac Total IgA performed initially</p>	<ul style="list-style-type: none"> • <u>Normal IgA, negative TTG</u> (IgA at or above Normal Low for patient's age, TTG IgA <=6.9) (Celiac disease unlikely. Further reflex testing not indicated.) • <u>Normal IgA, positive TTG</u> (IgA at or above Normal Low for patient's age, TTG IgA >=10.1) (Celiac disease likely. Further reflex testing not indicated.) • <u>Normal IgA, equivocal TTG</u> (IgA at or above Normal Low for patient's age, TTG IgA 7.0 - 10.0) (Endomysial Ab IgA ordered by reflex rule for further evaluation) • <u>Decreased IgA, negative TTG</u> (IgA below Normal Low for patient's age, TTG IgA <=6.9), (Transglutaminase Ab IgG, Gliadin Ab IgG, and Gliadin Ab IgA ordered by reflex rule for further evaluation.) • <u>Decreased IgA, positive TTG IgA</u> (IgA below Normal Low for patient's age, TTG IgA >=10.1) (Transglutaminase Ab IgG, Gliadin Ab IgG, and Gliadin Ab IgA ordered by reflex rule for further evaluation.) • <u>Decreased IgA, equivocal TTG IgA</u> (IgA below Normal Low for patient's age, TTG IgA 7.0 - 10.0): (Endomysial Ab IgA, Transglutaminase Ab IgG,

INITIAL TEST and RESULT	CONFIRMATION TESTING/ ADDITIONAL WORKUP
	Gliadin Ab IgG, and Gliadin Ab IgA ordered by reflex rules for further evaluation.
Cerebral Spinal Fluid (CSF) If Protein Electrophoresis [Test Code 8651] is requested	IgG Index Synthesis CSF Careset [Test Code 11593], IgG Index Serum [Test Code 11592]
Cerebral Spinal Fluid (CSF) RBC Cell Count greater than or equal to 25 cells in tube 3	Additional count of tube 1
Cerebral Spinal Fluid (CSF) WBC Cell Count greater than 0, in tube 3	Manual differential
Cryptococcus Antigen ordered on CSF	Cryptococcus CSF Panel (Cryptococcus Ag and Fungal Culture)
Clostridium difficile by EIA Toxin Positive Antigen Negative Toxin Negative Antigen Positive	Clostridium difficile by PCR
Culture from catheter tip or foreign bodies	Culture catheter/device
Direct Antiglobulin	ABO/RH
Mononucleosis Screen, Epstein Barr (EBV) IgM if Negative (Test Code 236): If Mononucleosis Screen is Negative	Epstein Barr (EBV) VCA IgM Acute Antibody (Test Code 4268)
All Hematopoietic neoplasms	DNA extraction and hold, cytogenetics, and flow cytometry
Hemoglobin (Hgb) A2 result is greater than 10%	Capillarys Hemoglobin Electrophoresis (to confirm Hgb E)
Hemoglobin Fractionation that identifies new Hemoglobin S	Sickle Cell Screen
Hemoglobin Fractionation	CBC on all orders for Hemoglobin Fractionation that do not already have a CBC ordered in the past 30 days.
Hepatitis C Genotype w/Amplification: Positive Hepatitis C Virus amplification result	Hepatitis C Genotype
Her2/neu immunohistochemistry 2+ positive	FISH for Her2/neu
Herpes Simplex, IgM, Antibody Screen result is positive or equivocal	Herpes Ab IgM IFA
HIV 1/2 Antibody EIA and Quick Test If HIV 1/2 Antibody is positive	<ul style="list-style-type: none"> a) HIV Quick test. If Positive: reflex to Roche HIV. b) Roche HIV: If positive for Antibody then reflex to Geenius for confirmation. (Positive Antigens do not reflex; HIV RNA testing requires a new order and a new specimen).
HIV Quick (Spectrum Health Regional Lab only)	Cancel and reorder as HIV ½ Antibody EIA
Immunohistochemical (IHC) Stain [Test Code 1098] All newly diagnosed patients with endometrial cancer, including cases of recurrence when no IHC was previously performed. <ul style="list-style-type: none"> a) If MLH1 is absent: b) If PMS2, MSH2, and/or MSH6 is/are absent: 	MLH1 Promoter Hypermethylation Analysis [Test Code 7091] If hypermethylation is absent: refer to genetics consult. Refer to genetics consult.

INITIAL TEST and RESULT	CONFIRMATION TESTING/ ADDITIONAL WORKUP
<p>Leukemia/Lymphoma/Myeloma and/or Non-Hodgkin lymphoma panels by flow cytometry: if indicated, reflex testing may be added to further characterize possible abnormal cell populations identified by the screening panel.</p> <p>These panels are reviewed continuously in multidisciplinary conferences and by the flow cytometry laboratory and hematopathologists.</p>	<p>The following add on panels may be employed after initial testing, as needed and appropriate, to further evaluate any possible abnormal population of cells.</p> <p>B lymphoblastic leukemia (B-ALL) panel T lymphoblastic leukemia (T-ALL) panel Chronic lymphocytic leukemia (CLL) panel Hairy cell leukemia (HCL) panel Extended B-cell tube panel Extended T-cell tube panel NK cell or LGL panel CD10 positive B-cell panel CD5 positive B-cell panel Acute myeloid leukemia (AML) panel Extended myeloid or monocytic panel Plasma cell panel Mast cell panel Solid tumor panel Neuroblastoma panel</p>
<p>NICU patients with a cord blood workup</p>	<p>Antibody Screen</p>
<p>Non SHGR 1) Patients with a diagnosis of Influenza during Influenza season 2) Patients with a diagnosis of Influenza during Non-influenza season</p>	<p>Influenza (Flu) A/B Rapid [test code 4626]; If Influenza (Flu) A/B Rapid is Negative, then Influenza (Flu) A/B PCR [test code 156]</p> <p>Influenza (Flu) A/B PCR [test code 156]</p>
<p>SHGR: Influenza A/B Rapid (Test Code 155); if negative</p>	<p>Influenza A/B PCR (Test Code 156)</p>
<p>Adults (pts >17) Inpatient or Observational at Butterworth or Blodgett with active warfarin order and no active INR order</p>	<p>LIS will automatically order a PT/INR when: The rule is evoked upon administration of a warfarin dose being signed in the eMAR. Once evoked the rule looks for:</p> <ol style="list-style-type: none"> 1. Pt age > 17 years, 2. Inpatient, Observation encounters at BW, BL if 1 and 2 are true then rule keeps evaluating for, 3. an active PT(with INR) order with a requested date and time within a future 24 hours of the administration of the warfarin dose. or 4. an active PT (with INR) with a frequency of; q 24hr, q am, q daily. (child order to be placed approx 00:30 with an ops job) <p>If neither 3 or 4 of these conditions are met the rule will order orderable Protime (with INR) w/ order entry fields.</p>
<p>Malaria Rapid Screen (Test Code: 7087) 1) If presumptive positive for malaria antigens 2) If presumptive negative for malaria antigens</p>	<ol style="list-style-type: none"> 1) Parasitemia Level (Test Code: 7089); Malaria speciation confirmation by thin/thick smear microscopy evaluation 2) Negative result confirmed by thin/thick smear microscopy evaluation. If smear is positive,

INITIAL TEST and RESULT	CONFIRMATION TESTING/ ADDITIONAL WORKUP
	then reflex to Parasitemia Level (Test Code: 7089)
Myeloproliferative Neoplasms (MPN) 1. JAK2 V617F ordered without prior 12 month BCR-ABL 2. JAK2 V617F mutation negative with BCR-ABL negative within past 12 months.	1.BCR-ABL must be performed first if Negative then JAK2 V617F performed 2.Then MPN Expanded Panel (Test Code 7090)
All newly diagnosed of MDS and MDS/MPN	Heme molecular Sequence Analysis
Pathologist Review (Test Code: 8367) If review of peripheral blood smear is ordered without required accompanying CBC with differential, and if the specimen is within 10 hours of collection.	Complete Blood Count (CBC) with Differential (Test Code: 8411)
Platelet Count less than 100,000/ μ L Updated November 2021	Immature platelet fraction (IPF)
Patient is less than 6 months of age and has suspected Hgb S by the Hgb fractionation test	Capillarys Hemoglobin Electrophoresis
Patient is suspected of having Hgb C by the Hgb fractionation test	Capillarys Hemoglobin Electrophoresis
Platelet Aggregation Studies	Platelet Count
Platelet Function Assay (PFA 100) with Collagen/Epinephrine results greater than 180 seconds.	Collagen/ADP test
Platelet Function Assay (PFA)	Platelet Count and hematocrit
Factor 8 Activity for patients on emicizumab (Hemlibra) therapy	Chromogenic Factor 8 Activity
Positive Antibody Screen, or a positive Direct Antiglobulin Test (DAT) on inpatients, outpatients, and surgical patients	Relevant studies as needed including antibody identification, antigen typing, direct antiglobulin test, elution and absorption. In addition, packed blood cells will be antigen typed and crossmatched.
Positive culture for pathogen or organism with clinically significant concentration (bacteria or mycobacteria)	Susceptibility and typing as necessary.
Positive Cryoglobulin test	Positive Cryoglobulins which have not had an identification in the past 2 months will have the Reflex Cryoglobulin Interpretation ordered.
Positive Group B Strep, Penicillin allergy, PCR on OB patients	Susceptibility testing.
Positive Hepatitis B surface antigen	HbsAg Confirmation test
Positive Lyme Disease Screen	Western Blot
Positive Amphetamine, Cannabinoids, Ethanol, Methadone, opiates, Oxycodone or cocaine on a Drug of Abuse screen for Inpatient Obstetric Patients (Mothers and their babies).	LC/MS Confirmation
Positive opiates on Comprehensive Drug Screens	LC/MS Confirmation
Positive Gamma HydroxyButyrate (GHB)	GC/MS Confirmation (Mayo)
Positive prenatal Profile Type &Antibody Screen	Antibody identification with titer if identified antibody is clinically significant
Prostate Specific Antigen (PSA) Free Level	PSA total on all orders that do not already have a PSA ordered on the same specimen.
Positive Syphilis IgG Antibody	RPR Titer at Spectrum Health, also confirmation by MHA-TPA at State Health Department
Rapid Plasma Reagin (RPR) test for syphilis	SYPHILIS IgG SCREEN will be performed instead of the RPR.
Prothrombin Time (PT) Mixing Study	PT and, if the PT Mixing Study does not correct, order a Lupus Anticoagulant Screen (LA1)
Random urine microalbumin	Urine creatinine
Sputum Culture without gram stain order	Gram stain for specimen suitability. If unsuitable, "culture" is cancelled.
Sputum Gram Stain without culture	Sputum Culture ordered. If Gram Stain is

INITIAL TEST and RESULT	CONFIRMATION TESTING/ ADDITIONAL WORKUP
	unsuitable, "culture" is cancelled.
Thyroglobulin	Automatically cancel AntiTgAB request when ordered with srTg
Type & Screen (T&S) on a patient with autologous or directed units	Crossmatch of the units
Type & Screen (T&S) on a pre-op patient with an antibody	Crossmatch of two antigen negative units
Metastatic Gastric and Esophageal Adenocarcinomas	Perform Her2/neu immunohistochemistry. When metastasis is pathologically confirmed (can use either metastatic site or primary tumor). <ul style="list-style-type: none"> • Equivocal immuno results will reflex to FISH testing.
All colorectal carcinoma resections at Butterworth and Blodgett Hospitals	<ul style="list-style-type: none"> • Immunohistochemical (IHC) protein studies for MSH2, MSH6, MLH1, PMS2 [Test Code 1098] • PCR for Microsatellite Instability (MSI)[Test Code 904] • Additional reflex to BRAF [Test Code 709], if MSI-H with loss of MLH1 on IHC
Metastatic Colorectal Carcinoma	Colon Mutation Analysis Panel (Test Code 7094) -includes genotyping for KRAS, NRAS, and BRAF mutations
All Diffuse Large B-cell Lymphoma/other B-cell Lymphoma with aggressive or high grade features	<ol style="list-style-type: none"> 1. Immunohistochemical/In situ hybridization staining which may include: CD3, CD20, CD5, CD10, BCL-1, Ki-67, EBERish, BCL-2, BCL-6, MUM1, MYC, CD30, and CD45 2. Cytogenetic FISH testing to include: BCL-2, BCL-6, and MYC rearrangement testing, which will reflex to t(8;14) when appropriate per FISH lab protocol.
All newly diagnosed lung (non-small cell) adenocarcinomas, stage 4 only	Molecular Lung Cancer Panel (Test Code 7030)
Metastatic melanoma Including Cases with Positive lymph nodes	BRAF IHC if IHC negative reflex to BRAF molecular testing
Glioblastoma (brain) Grade 4 Anaplastic Astrocytoma IDH-wildtype	MGMT Methylation Analysis [Test Code706] p53 [Test Code1098] IDH1 [Test Code 1098] Cancer Hotspot analysis
Glioblastoma (brain) Grade 4 Recurrent Glioblastoma initially MGMT positive (Methylated)	MGMT Methylation Analysis [Test Code706]
Glioma Grade 2 and 3	p53 [Test Code 1098] IDH1 [Test Code 1098] Del 1p/19q by FISH [Test Code 955] Ki-67 [Test Code 1098] ATRX by Immunohistochemistry
All newly diagnosed or previously untested oropharyngeal squamous cell carcinomas	p16 immunohistochemistry (surrogate marker for HPV)
Pain Management Panel If positive screen result:	<ul style="list-style-type: none"> ○ Amphetamines-reflex to methamphetamine and amphetamine if screen positive ○ Barbiturates-reflex to Butalbital, Secobarbital, and Pentobarbital if screen positive ○ Benzodiazapines-reflex to Benzodiazapines Urine Level Panel if screen positive ○ Cannabinoids-reflex to confirmation if positive ○ Cocaine Metabolite-reflex to confirmation if screen positive ○ Ethanol-reflex to confirmation if screen

INITIAL TEST and RESULT	CONFIRMATION TESTING/ ADDITIONAL WORKUP
	positive <ul style="list-style-type: none"> ○ Methadone Metabolite-reflex to Methadone and EDDP if screen positive
UA or UA culture if (where volume is inadequate for microscopic exam)	Urine Dipstick (U dip)
Antibody Titer	ABO/Rh and Antibody Screen
Tissue Specimens ordered as a Body Fluid Culture	Cancel and Order as a Tissue Culture
Body Fluid Specimens ordered as a Tissue Culture	Cancel and Order as a Body Fluid Culture
Specimens in Abbott Multi Collect tube ordered as Aptima specimens	Cancel Aptima test and reorder corresponding Abbott test for Chlamydia PCR, Gonococcus PCR
Specimens in Aptima collection tube ordered as Abbott Multi Collect tube specimens	Cancel Abbott test and reorder corresponding Aptima test for Chlamydia NAAT, Gonococcus NAAT, or Trichomonas Only
Lupus Screen LA1 elevated	LA2 if elevated perform Protime and aPTT if these are elevated perform mixing study for Protime/aPTT Pathologist interpretation reported with elevated results and/or mixing study
Heparin Dependent Antibody (HIT) Positive or Borderline	Serotonin Release Assay (SRA)
Platelet Aggregation	Hematocrit
All newly diagnosed adult patients older than 80 with newly diagnosed AML	IDH Mutation Analysis
Bone marrow or blood EDTA samples (Philadelphia positive B-ALL and CML)	RNA extract and hold
Mycoplasma pneumoniae IgM reactive or equivocal	Mycoplasma pneumoniae IgM by IFA
Blastomyces Antibody by EIA, Serum equivocal or Positive	Blastomyces Antibody by Immunodiffusion
Serum Protein Electrophoresis IFE if indicated	Serum Protein Electrophoresis with paraprotein found or elevation of protein fraction in electrophoresis pattern. Reflex to Immunofixation for identification.
24 hour Urine Protein Electrophoresis IFE if indicated	Urine Protein Electrophoresis with paraprotein found or elevation of protein fraction in electrophoresis pattern. Reflex to Immunofixation for identification.
Random Urine Protein Electrophoresis IFE if indicated	Urine Protein Electrophoresis with paraprotein found or elevation of protein fraction in electrophoresis pattern. Reflex to Immunofixation for identification.
HSV viral culture from cutaneous and mucocutaneous lesions	Cancel and change to HSV PCR order
Group A Streptococcus negative antigen test on pediatric patients	Throat Culture
Pap test and HPV	Reflex Information Is HPV Requested? a. If NO HPV testing is desired, select NO and only the pap test will be ordered. b. If YES, HPV testing is desired, select one of two options: i. CO-TESTING (30-64 y/o) 1. If Co-testing is selected, the HPV test will be ordered and performed regardless of the pap test final diagnosis. 2. Note: Co-testing is recommended for patients age 30-64. ii. HPV REFLEX (see link below for criteria) 1. If reflex is selected the HPV test will only be performed in the following scenarios:

INITIAL TEST and RESULT	CONFIRMATION TESTING/ ADDITIONAL WORKUP
	<p>a. The pap test final diagnosis is NIL and the patient is between ages 30-64.</p> <p>b. The pap test final diagnosis is ASCUS and the patient is between ages 21-64.</p> <p>c. The pap test final diagnosis is LSIL and the patient is between ages 25-64 and is not pregnant.</p> <p>c. The HPV Genotype test will automatically reflex for all patients between the ages of 30-64 with a NIL pap test diagnosis and a positive HPV test diagnosis.</p> <p>1. Note: Add on HPV and HPV Genotype tests can be added up to 4 weeks from the collection date</p>
Body Fluid culture greater than 1mL sample with only aerobic culture	Add anaerobic culture
GYN sentinel lymph nodes which show no evidence of metastasis on initial H&E levels (applies to all GYN sentinel lymph nodes, including, but not limited to, those removed for endometrial, vulvar, and cervical carcinomas).	Cytokeratin AE1/AE3 (to be performed on all blocks of all nodes deemed negative for metastasis following initial review of H&E sections).
<p>Testing for RHD genotyping (molecular testing) to further determine patient management as RHD positive or negative. Additional ability to order molecular testing for patients with difficult antibody situation, as deemed necessary by the Blood Bank physician/pathologist.</p> <p>For more details please see original MEC from 8/2021</p>	

Optional Reflex Testing

ORDER	INITIAL TEST and RESULT	OPTIONAL FOLLOW UP TESTING
ANA Screen	ANA EIA if positive	ANA Hep2 (IFA) if positive reflex to titer. If titer is greater than 1:160 reflex to Anti-dsDNA, anti-Sm, anti-RNP, anti-SSA, anti-SSB, anti Scl70, anti-centromere and anti-Jo1
Breast Biopsy	Primary and recurrent/metastatic Invasive Mammary Carcinoma (excisional biopsy / lumpectomy / mastectomy)	Immunohistochemistry for estrogen receptor, progesterone receptor, Her2/neu protein
Breast Biopsy	Ductal Carcinoma In Situ	Immunohistochemistry stains for estrogen receptor, progesterone
CBC w/diff	WBC less than 3.0 or greater than 18.0 HGB less than 8.0 or greater than 18.0 MCV less than 75.0 or greater than 105.0 Absolute neut count less than 1.50 or greater than 9.00 Absolute lymph count less than 0.39 or greater than 4.50 Absolute mono count greater than 1.50 Absolute eos count greater than 1.00 Absolute bas count greater than 0.20 Abnormal instrument flags suggesting abnormality	Manual WBC differential
CBC	CBC specimens that fulfill criteria listed in Pathologist Review (Hematology #6783)	Pathologist review
CBC w/ Diff	CBC specimens that fulfill criteria listed in Pathologist Review (Hematology #6783)	Pathologist Review
Cell Ct only BFL	Body fluid specimens that fulfill criteria listed in Pathologist Review (Hematology #6783)	Pathologist review
Fetal Cells by Flow Cytometry	Ordered STAT and received in lab outside of flow cytometry testing hours (after 3:30 Mon-Fri or after 10:00am Sat or anytime Sunday)	Fetal Hemoglobin by Kleihauer Betke performed in place of flow cytometry test
Triglyceride do LDL Direct if >400	LDL Direct if the triglycerides are greater than 400.	LDL Direct
Lipid Panel do LDL Direct if Triglycerides >400	LDL Direct if the triglycerides are greater than 400.	LDL Direct
Pap do HPV if	Cervical Cytology with ASCUS, ASC-H OR LSIL	HPV-high risk
PAP Test - HPV if ASCUS or AGUS	Cervical Cytology with ASCUS or AGUS	HPV
PAP Test - HPV if NIL, ASCUS or AGUS	Cervical Cytology with NIL, ASCUS or AGUS	HPV
Preg Serum Quant Progesterone if	HCG greater than 5 mIU/mL	Progesterone level
PSA Sym FPSA if	PSA between 2.5 and 10 ng/mL	Free PSA
Strep A, PCR if Negative	Negative rapid Strep A	Throat Culture
TSH FT4 if indicated	TSH less than 0.3 OR greater than 5.0 mCU/mL	Free T4
Biopsy of Thyroid Aspirate	Diagnosis of atypical follicular neoplasm of uncertain significance and suspicious	Afirma GEC (Veracyte)

ORDER	INITIAL TEST and RESULT	OPTIONAL FOLLOW UP TESTING
	for follicular/hurthle cell neoplasm (Bethesda Classification-Categories 3 and 4)	
Thyroid Function Cascade	<ol style="list-style-type: none"> 1. TSH above 5.0 2. TSH below 0.3 <ol style="list-style-type: none"> a. TSH below 0.1 and FT4 below 1.6 	<ol style="list-style-type: none"> 1. Reflex to FT4 and TPO if TSH is high (above 5.0). 2. Reflex to FT4 if TSH is low (below 0.3). <ol style="list-style-type: none"> a. Reflex to Free T3 if TSH is very low (below 0.1) and FT4 is normal (below 1.6)
UA culture if	Urinalysis (UA) with two or more of the following abnormal findings, provided there are less than 10 squamous epithelial cells observed per high power field: -Greater than or equal to 10 WBC -Positive leukocyte esterase -Positive nitrite OR if specimen is -Grossly bloody	Culture and sensitivity
UA culture if (Where volume is inadequate for microscopic exam)	Volume is inadequate for microscopic exam and Urine Dipstick (U dip) with one or more of the following abnormal findings: -Positive leukocyte esterase -Positive nitrite	Culture and sensitivity
Hepatitis C Virus Antibody do HCV RNA if indicated	Positive or Indeterminate Hepatitis C Virus Antibody result	HCV RNA
Leukemia or Non-Hodgkin Lymphoma Panel by Flow Cytometry	Cell population is diagnostic of circulating leukemia/lymphoma/myeloma Patients under age of 80 with new diagnosis.	FISH testing
Peanut IgE Reflex	Peanut IgE => 0.10 kU/L	Peanut component allergen panel (Peanut Ara h 1 IgE, Peanut Ara h 2 IgE, Peanut Ara h 3 IgE, Peanut Ara h 8 IgE, and Peanut Ara h 9 IgE)
Egg IgE Reflex	Egg white IgE =>0.10 kU/L	Egg component allergen Panel (Ovomucoid IgE and Ovalbumin IgE)
Milk IgE Reflex	Milk (cow) =>0.10 kU/L	Milk component allergen Panel (Casein IgE, Alpha-Lactalbumin IgE, and Beta-lactoglobulin IgE)